

## Contents Volume 13, 1988/1989

|   |     |
|---|-----|
| Palaeoceanographic record of the last deglaciation in the Strait of Sicily<br>C. Vernaud-Grazzini (Paris, France), A.M. Borsetti (Bologna, Italy), F. Cati (Bologna, Italy), P. Colantini (Bologna, Italy), S.D'Onofrio (Bologna, Italy), J.F. Saliège (Paris, France), R. Sartori (Bologna, Italy) and R. Tampieri (Torino, Italy) .....   | 1   |
| Palaeoceanography of the mid-Pleistocene<br>S. Gartner (College Station, TX, U.S.A.) .....  | 23  |
| Differential dissolution of Upper Cretaceous deep-sea benthonic foraminifers from the Angola Basin, South Atlantic Ocean<br>J.G.V. Widmark and B.A. Malmgren (Uppsala, Sweden) .....  | 47  |
| Diatom productivity peak and increased circulation during latest Quaternary: Alboran Basin (Western Mediterranean)<br>F. Abrantes (Narragansett, RI, U.S.A.) .....  | 79  |
| Bathymetric distribution of benthic foraminifera on the Australian-Irian Jaya continental margin, eastern Indonesia<br>L.J. Van Marle (Amsterdam, The Netherlands) .....  | 97  |
| Dinoflagellate cysts, sea level changes and planktonic foraminifers across the Cretaceous-Tertiary boundary at El Haria, northwest Tunisia<br>H. Brinkhuis and W.J. Zachariasse (Utrecht, The Netherlands) .....  | 153 |
| Neogene stratigraphy, foraminifera, diatoms, and depositional history of Maria Madre Island, Mexico: evidence of Early Neogene marine conditions in the southern Gulf of California<br>C. McCloy (Stanford, CA, U.S.A.), J.C. Ingle (Stanford, CA, U.S.A.) and J.A. Barron (Menlo Park, CA, U.S.A.) .....   | 193 |
| Late Holocene deep-sea benthic foraminifera from the Sea of Marmara<br>S.N. Alavi (Erdemli, Turkey) .....   | 213 |
| Extinction, survivorship and evolution of planktic foraminifera across the Cretaceous/Tertiary boundary at El Kef, Tunisia<br>G. Keller (Princeton, NJ, U.S.A.) .....   | 239 |
| Late glacial to recent deep-sea benthic foraminifera from the northeastern Atlantic (Cadiz Gulf) and western Mediterranean (Alboran Sea): paleoceanographic results<br>M.-H. Caralp (Talence, France) .....   | 265 |
| Cenozoic Cosmopolitan Deep-Water Benthic Foraminifera by F.P.C.M. Van Morkhoven, W.A. Berggren and A.S. Edwards — J.H. Lipps .....  | 291 |
| Conodonts Investigative Techniques and Applications by R.L. Austin — K.J. Müller .....  | 291 |
| Radiolarian-based transfer-functions for estimating paleoceanographic conditions in the south Indian Ocean<br>J.J. Morley (Palisades, NY, U.S.A.) .....   | 293 |
| Latitude-related shell patterns in Radiolaria: <i>Botryostrobus auritus/australis</i> morphotypes in the equatorial to arctic Pacific<br>D. Boltovskoy and Vrba (Buenos Aires, Argentina) .....   | 309 |
| Palaeoceanographic interpretations of coccoliths and oxygen-isotopes from the sediment surface of the southwest Indian Ocean<br>M.J. Fincham and A. Winter (Rondebosch, South Africa) .....   | 325 |
| Quaternary surface-water stable isotope from calcareous nanofossils at DSDP Site 593, southern Tasman Sea<br>W.C. Dudley (Hilo, HI, U.S.A.) and C.S. Nelson (Hamilton, New Zealand) .....   | 353 |
| Paléocéologie des associations d'ostracodes d'une formation carbonatée à Rudistes en environnement péri-deltaïque: l'exemple du Santonien de la Cadière d'Azur (Sud-Est France). [Paleoecology of ostracode assemblages in a carbonate bank with rudists (perideltaic context): example of the Santonian of La Cadière d'Azur (southeast France)]<br>D. Grosheny and J.F. Babinot (Marseille, France) ..... | 375 |